

BOGOLIUBOV SPECTRUM OF INTERACTING BOSE GASES

MATHIEU LEWIN, PHAN THÀNH NAM, SYLVIA SERFATY,
AND JAN PHILIP SOLOVEJ

ABSTRACT. We study the large- N limit of a system of N bosons interacting with a potential of intensity $1/N$. When the ground state energy is to the first order given by Hartree's theory, we study the next order, predicted by Bogoliubov's theory. We show the convergence of the lower eigenvalues and eigenfunctions towards that of the Bogoliubov Hamiltonian (up to a convenient unitary transform). We also prove the convergence of the free energy when the system is sufficiently trapped. Our results are valid in an abstract setting, our main assumptions being that the Hartree ground state is unique and non-degenerate, and that there is complete Bose-Einstein condensation on this state. Using our method we then treat two applications: atoms with "bosonic" electrons on one hand, and trapped 2D and 3D Coulomb gases on the other hand.

CONTENTS

1. Introduction	2
2. Main abstract results	5
2.1. Assumptions	5
2.2. The Bogoliubov Hamiltonian	8
2.3. Exciting the Hartree state	10
2.4. Convergence of the excitation spectrum	11
2.5. Positive temperature	15
3. Applications	16
3.1. Bosonic atoms	16
3.2. Trapped Coulomb gases	20
4. Operators on Fock spaces	24
5. Bound on truncated Fock space	27
6. Localization in Fock space	32
7. Proof of Main Theorems	33
7.1. Proof of Theorem 2	33
7.2. Proof of Theorem 3	37
Appendix A. Bogoliubov Hamiltonian	40
Appendix B. Localization of band operators on \mathcal{F}_+	46
Appendix C. Logarithmic Lieb-Oxford inequality	46
References	49

Date: April 24, 2013. Final version to appear in *Comm. Pure Appl. Math.*

